

Appendix

Syntheses methods

Unit-of analysis issues were handled as follows: In studies reporting on more than two treatment groups, three approaches were taken to avoid unit-of-analysis conflicts: In case of groups being comparable, we combined them. If additional groups used treatments not in accordance with current standard (e.g. multiple-step treatment without any root-canal medication), this group was omitted. If a factorial design was used (e.g. both groups were compared in vital and non-vital teeth), with separate reporting for all groups, we compared subgroups and handled them as if they were separate studies for meta-analysis.

Meta-regression was additionally performed. As some studies did not clearly state randomization (see above), a sensitivity analysis excluding these studies was performed. Similarly, as some studies reported results to have not been significantly different (but did not report on exact effect estimates), we imputed the number of events per group as the mean event rate in a sensitivity analysis, making best use of all available information. For subgroup comparisons, Chi-square test was performed. For meta-regression, the unrestricted maximum-likelihood method was used; Bonferroni adjustment to correct for multiple testing was planned, but not required, as no significant associations were found even without such correction.

Trial sequential analysis was performed. RIS was calculated based on type I error risk of $\alpha=0.05$ and a type II error risk of $\beta=0.20$ (equivalent to a power of 0.80). The control event proportion (i.e. event incidence in multiple-visit group) and the relative risk reduction (RRR) were used to estimate RIS. RRR was based on an a priori defined worthwhile interventional effect of 20% (lower effects might be worthwhile, but would increase RIS even further) (1, 2). RIS was diversity (heterogeneity) adjusted (DARIS). To assess if differences yielded by conventional meta-analysis are robust, TSA additionally estimates trial sequential monitoring boundaries (TSMB), i.e. statistical thresholds for significance which are adapted depending on the so far reached sample size. The Lan-DeMets version (3) of the O'Brien–Fleming

function (4) was used for calculating the TSMBs. In case the cumulative Z-value crossed the conventional boundary of significance ($Z=\pm 1.96$) but not the TSMBs for benefit or harm, we defined such findings as spuriously significant. Firm evidence was assumed to be reached when the Z-curve crossed the TSMB for benefit or harm before the DARIS was reached. Effect estimates supported by only few small trials are thus handled stricter than those supported by large samples. In addition to such superiority/inferiority TSMBs, monitoring boundaries for futility were calculated (these indicate if further trial conduct is likely to be futile, i.e. if sufficient evidence has been accrued to claim non-inferiority of treatments). Further details regarding the applied method to calculate TSMB have been reported elsewhere (1). TSA was performed with TSA 0.9 (Copenhagen Trial Unit, Copenhagen, Denmark) (5).

Table S1: Excluded Studies

Soltanoff 1978 (6)	Selection bias (allocation according to tooth status)
O'Keefe 1976 (7)	Selection bias (allocation according to available time)
EIMubarak 2010 (8)	No RCT
Raju 2014 (9)	Did not compare 1- vs 2 visits
Xavier 2013 (10)	No clinical outcome
Bhagwat 2013 (11)	Did not compare 1- vs 2 visits
Roane 1983 (12)	No RCT
Oliet 1983 (13)	Selection bias (allocation according to patient acceptance, available time, symptoms of tooth)
Ether 1978 (14)	Not available
Eleazer 1998 (15)	no RCT
Fava 1989 (16)	Compared different techniques
Fox 1970 (17)	Did not compare 1- vs 2 visits
Genet 1986 (18)	Did not compare 1- vs 2 visits
Morse 1987 (19)	Did not compare 1- vs 2 visits
Yesilsoy 1988 (20)	Did not compare 1- vs 2 visits
Trope 1991 (21)	Did not compare 1- vs 2 visits
Koba 1999 (22)	Did not compare 1- vs 2 visits
Glennon 2004 (23)	Did not compare 1- vs 2 visits
Ng 2004 (24)	no RCT
Georgopoulou 1986 (25)	no RCT
Jurcak 1993 (26)	no RCT
Imura 1995 (27)	no RCT
Walton 1992 (28)	no RCT
Alacam 1985 (29)	Did not compare 1- vs 2 visits
Torabinejad 1994 (30)	Did not compare 1- vs 2 visits
Sjögren 1990 (31)	Did not compare 1- vs 2 visits
Siqueira 2002 (32)	Did not compare 1- vs 2 visits
Orstavik 1996 (33)	Did not compare 1- vs 2 visits
Perkruhn 1986 (34)	Did not compare 1- vs 2 visits
Kvist 2004 (35)	no clinical outcomes reported (CFU)
Rudner 1981 (36)	no RCT
Kenrick 1999 (37)	no RCT
Sjögren 1997 (38)	Did not compare 1- vs 2 visits
Maddox 1977 (39)	Did not compare 1- vs 2 visits
Sjögren 1990 (31)	no RCT
Fleming 2010 (40)	no RCT
Singla 2008 (41)	pulpectomy
Kalhor 2009 (42)	no RCT
Shaikh 2013 (43)	Not available

Table S2: Risk of bias of included studies. Bias assessment followed guidelines outline by The Cochrane Collaboration (44).

	Sequence generation	Allocation Concealment	Blinding of operator	Blinding of examiner	Incomplete data	Selective reporting	Overall risk of bias
Akbar 2013 (45)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Albashaireh & Alnegreshi 1998 (46)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Al-Negrish & Habahbeh 2006 (47)	Unclear	Unclear	High	High	Low	Low	Unclear/High
DiRenzo 2002 (48)	Low	Low	Low	Low	Low	Low	Low
Dorsani 2013 (49)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Fava 1989 (50)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Fava 1994 (51)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Gesi 2006 (52)	Low	Low	High	Low	Low	Low	Unclear/High
Ghoddusi 2006 (53)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Ince 2009 (54)	Unclear	Unclear	Unclear	Low	Low	Low	Unclear/High
Jabeen 2014 (55)	Low	Low	Low	Low	Low	Low	Low
Liu & Leng 2013 (56)	Unclear	Unclear	High	Low	Low	Low	Unclear/High
Molander 2007 (57)	High	Unclear	High	Low	Low	Low	Unclear/High
Mulhern 1982 (58)	Low	Low	High	Low	Low	Low	Unclear/High
Oginni 2004 (59)	Low	Low	Low	Low	Low	Low	Low
Paredes-Vieyra 2012 (60)	Unclear	High	High	Low	Low	Low	Unclear/High
Pekruhn 1981 (61)	Unclear	Unclear	Unclear	Low	Low	Low	Unclear/High
Penesis 2008 (62)	Low	Low	Low	Low	Low	Low	Low
Peters and Wesslink 2002 (63)	Low	Unclear	High	Low	Low	Low	Unclear/High
Prashanth 2011 (64)	Unclear	High	High	High	High	Low	Unclear/High
Rao 2014 (65)	Unclear	High	Low	Low	Low	Low	Unclear/High
Risso 2008 (66)	Low	Low	Low	Low	Low	Low	Low
Singh and Kargh 2012 (67)	Unclear	High	Low	Low	Low	Low	Unclear/High
Trope 1999 (68), Walimo 2005 (69)	Low	Low	Low	Low	Low	Low	Low
Wang 2010 (70)	Low	Low	High	High	Low	Low	Unclear/High
Weiger 2000 (71)	Unclear	High	High	Low	Low	Low	Unclear/High
Wong 2015a (72)	High	High	Low	Low	Low	High	Unclear/High
Wong 2015b (73)	Low	Unclear	High	High	Low	Low	Unclear/High
Yoldas 2004 (74)	Unclear	Unclear	High	Low	Low	Low	Unclear/High




Figure S1: Funnel plots. (a) Risk long-term complications, (b) risk of experiencing any postoperative pain, (c) risk of experiencing a flare-up. Standard errors are plotted against logRR to estimate possible small study effects or publication bias via an asymmetry of the funnel. White circles: estimates reported by included studies, black balls: imputed estimates in case of suspected publication bias. White diamond: effect estimate based on included studies, black diamond: effect estimate based on included and imputed studies.

References for the appendix

1. Schwendicke F, Goestemeyer G, Gluud C. Cavity lining after excavating caries lesions: meta-analysis and trial sequential analysis of randomized clinical trials. *Journal of Dentistry* 2015;doi: **10.1016/j.jdent.2015.07.017**.
2. Schwendicke F, Karimbux N, Allareddy V, Gluud C. Periodontal treatment for preventing adverse pregnancy outcomes: a meta- and trial sequential analysis. *PloS one* 2015;**10**(6):e0129060.
3. Demets DL, Lan KKG. Interim analysis: The alpha spending function approach. *Statistics in medicine* 1994;**13**(13-14):1341-52.
4. O'Brien PC, Fleming TR. A Multiple Testing Procedure for Clinical Trials. *Biometrics* 1979;**35**(3):549-56.
5. Thorlund K, Engstrøm J, Wetterslev J, Brok J, Imberger G, Gluud C. User Manual for Trial Sequential Analysis (TSA). http://www.ctu.dk/tsa/files/tsa_manual.pdf 2011:1-115.
6. Soltanoff W. A comparative study of the single-visit and the multiple-visit edodontic procedure. *Journal of endodontics* 1978;**4**(9):278-81.
7. O'Keefe EM. Pain in endodontic therapy: preliminary study. *Journal of endodontics* 1976;**2**(10):315-9.
8. El Mubarak AH, Abu-bakr NH, Ibrahim YE. Postoperative pain in multiple-visit and single-visit root canal treatment. *Journal of endodontics* 2010;**36**(1):36-9.
9. Raju TB, Seshadri A, Vamsipavani B, Abhilash K, Subhash AV, Kumari KV. Evaluation of pain in single and multi rooted teeth treated in single visit endodontic therapy. *Journal of international oral health : JIOH* 2014;**6**(1):27-32.
10. Xavier AC, Martinho FC, Chung A, Oliveira LD, Jorge AO, Valera MC, et al. One-visit versus two-visit root canal treatment: effectiveness in the removal of endotoxins and cultivable bacteria. *Journal of endodontics* 2013;**39**(8):959-64.
11. Bhagwat S, Mehta D. Incidence of post-operative pain following single visit endodontics in vital and non-vital teeth: An in vivo study. *Contemporary clinical dentistry* 2013;**4**(3):295-302.
12. Roane JB, Dryden JA, Grimes EW. Incidence of postoperative pain after single- and multiple-visit endodontic procedures. *Oral surgery, oral medicine, and oral pathology* 1983;**55**(1):68-72.
13. Oliet S. Single-visit endodontics: a clinical study. *Journal of endodontics* 1983;**9**(4):147-52.
14. al. ESe. A comparison of one and two visit endodontics. *J Farmacia Odontol* 1978(8):215.
15. Eleazer PD, Eleazer KR. Flare-up rate in pulpally necrotic molars in one-visit versus two-visit endodontic treatment. *Journal of endodontics* 1998;**24**(9):614-6.
16. Fava LR. One-appointment root canal treatment: incidence of postoperative pain using a modified double-flared technique. *International endodontic journal* 1991;**24**(5):258-62.
17. Fox J, Atkinson JS, Dinin AP, Greenfield E, Hechtman E, Reeman CA, et al. Incidence of pain following one-visit endodontic treatment. *Oral surgery, oral medicine, and oral pathology* 1970;**30**(1):123-30.
18. Genet JM, Wesselink PR, Thoden van Velzen SK. The incidence of preoperative and postoperative pain in endodontic therapy. *International endodontic journal* 1986;**19**(5):221-9.
19. Morse DR, Furst ML, Belott RM, Lefkowitz RD, Spritzer IB, Sideman BH. Infectious flare-ups and serious sequelae following endodontic treatment: a prospective randomized

trial on efficacy of antibiotic prophylaxis in cases of asymptomatic pulpal-periapical lesions. *Oral surgery, oral medicine, and oral pathology* 1987;**64**(1):96-109.

20. Yesilsoy C, Koren LZ, Morse DR, Rankow H, Bolanos OR, Furst ML. Post-endodontic obturation pain: a comparative evaluation. *Quintessence international* 1988;**19**(6):431-8.

21. Trope M. Flare-up rate of single-visit endodontics. *International endodontic journal* 1991;**24**(1):24-6.

22. Koba K, Kimura Y, Matsumoto K, Watanabe H, Shinoki T, Kojy R, et al. Post-operative symptoms and healing after endodontic treatment of infected teeth using pulsed Nd:YAG laser. *Endodontics & dental traumatology* 1999;**15**(2):68-72.

23. Glennon JP, Ng YL, Setchell DJ, Gulabivala K. Prevalence of and factors affecting postpreparation pain in patients undergoing two-visit root canal treatment. *International endodontic journal* 2004;**37**(1):29-37.

24. Ng YL, Glennon JP, Setchell DJ, Gulabivala K. Prevalence of and factors affecting post-obturation pain in patients undergoing root canal treatment. *International endodontic journal* 2004;**37**(6):381-91.

25. Georgopoulou M, Anastassiadis P, Sykaras S. Pain after chemomechanical preparation. *International endodontic journal* 1986;**19**(6):309-14.

26. Jurcak JJ, Bellizzi R, Loushine RJ. Successful single-visit endodontics during Operation Desert Shield. *Journal of endodontics* 1993;**19**(8):412-3.

27. Imura N, Zuolo ML. Factors associated with endodontic flare-ups: a prospective study. *International endodontic journal* 1995;**28**(5):261-5.

28. Walton R, Fouad A. Endodontic interappointment flare-ups: a prospective study of incidence and related factors. *Journal of endodontics* 1992;**18**(4):172-7.

29. Alacam T. Incidence of postoperative pain following the use of different sealers in immediate root canal filling. *Journal of endodontics* 1985;**11**(3):135-7.

30. Torabinejad M, Cymerman JJ, Frankson M, Lemon RR, Maggio JD, Schilder H. Effectiveness of various medications on postoperative pain following complete instrumentation. *Journal of endodontics* 1994;**20**(7):345-54.

31. Sjogren U, Hagglund B, Sundqvist G, Wing K. Factors affecting the long-term results of endodontic treatment. *Journal of endodontics* 1990;**16**(10):498-504.

32. Siqueira JF, Jr., Rocas IN, Favieri A, Machado AG, Gahyva SM, Oliveira JC, et al. Incidence of postoperative pain after intracanal procedures based on an antimicrobial strategy. *Journal of endodontics* 2002;**28**(6):457-60.

33. Orstavik D. Time-course and risk analyses of the development and healing of chronic apical periodontitis in man. *International endodontic journal* 1996;**29**(3):150-5.

34. Pekruhn RB. The incidence of failure following single-visit endodontic therapy. *Journal of endodontics* 1986;**12**(2):68-72.

35. Kvist T, Molander A, Dahlen G, Reit C. Microbiological evaluation of one- and two-visit endodontic treatment of teeth with apical periodontitis: a randomized, clinical trial. *Journal of endodontics* 2004;**30**(8):572-6.

36. Rudner WL, Oliet S. Single-visit endodontics: a concept and a clinical study. *Compendium of continuing education in dentistry (Jamesburg, N.J. : 1995)* 1981;**2**(2):63-8.

37. Kenrick S. Endodontics: a multiple-visit or single-visit approach. *Australian endodontic journal : the journal of the Australian Society of Endodontology Inc* 2000;**26**(2):82-5.

38. Sjogren U, Figdor D, Persson S, Sundqvist G. Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *International endodontic journal* 1997;**30**(5):297-306.

39. Maddox D, Walton R, Davis C. Incidence of posttreatment endodontic pain related to medication and other factors. *Journal of endodontics* 1977;**3**(12):447-52.
40. Fleming CH, Litaker MS, Alley LW, Eleazer PD. Comparison of classic endodontic techniques versus contemporary techniques on endodontic treatment success. *Journal of endodontics* 2010;**36**(3):414-8.
41. Singla R, Marwah N, Dutta S. Single Visit versus Multiple Visit Root Canal Therapy. *International journal of clinical pediatric dentistry* 2008;**1**(1):17-24.
42. Kalhoro FA, Mirza AJ. A study of flare-ups following single-visit root canal treatment in endodontic patients. *Journal of the College of Physicians and Surgeons--Pakistan : JCPSP* 2009;**19**(7):410-2.
43. Shaikh AG, Kalhoro FA, Dall AQ. Evaluation of post operative pain in single and multi visit endodontic therapy with rotary instruments. *Medical Forum Monthly* 2013;**24**(1):49-54.
44. Higgins JPT, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions. Version 5.10 (updated March 2011): The Cochrane Collaboration; 2011.
45. Akbar I, Iqbal A, Al-Omiri MK. Flare-up rate in molars with periapical radiolucency in one-visit vs two-visit endodontic treatment. *The journal of contemporary dental practice* 2013;**14**(3):414-8.
46. Albashaireh ZS, Alnegrish AS. Postobturation pain after single- and multiple-visit endodontic therapy. A prospective study. *Journal of dentistry* 1998;**26**(3):227-32.
47. Al-Negrish AR, Hababbeh R. Flare up rate related to root canal treatment of asymptomatic pulpally necrotic central incisor teeth in patients attending a military hospital. *Journal of dentistry* 2006;**34**(9):635-40.
48. DiRenzo A, Gresla T, Johnson BR, Rogers M, Tucker D, BeGole EA. Postoperative pain after 1- and 2-visit root canal therapy. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics* 2002;**93**(5):605-10.
49. Dorasani G, Madhusudhana K, Chinni SK. Clinical and radiographic evaluation of single-visit and multi-visit endodontic treatment of teeth with periapical pathology: An in vivo study. *Journal of conservative dentistry : JCD* 2013;**16**(6):484-8.
50. Fava LR. A comparison of one versus two appointment endodontic therapy in teeth with non-vital pulps. *International endodontic journal* 1989;**22**(4):179-83.
51. Fava LR. A clinical evaluation of one and two-appointment root canal therapy using calcium hydroxide. *International endodontic journal* 1994;**27**(1):47-51.
52. Gesi A, Hakeberg M, Warfvinge J, Bergenholtz G. Incidence of periapical lesions and clinical symptoms after pulpectomy--a clinical and radiographic evaluation of 1- versus 2-session treatment. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics* 2006;**101**(3):379-88.
53. Ghoddusi J, Javidi M, Zarrabi MH, Bagheri H. Flare-ups incidence and severity after using calcium hydroxide as intracanal dressing. *The New York state dental journal* 2006;**72**(4):24-8.
54. Ince B, Ercan E, Dalli M, Dulgergil CT, Zorba YO, Colak H. Incidence of postoperative pain after single- and multi-visit endodontic treatment in teeth with vital and non-vital pulp. *European journal of dentistry* 2009;**3**(4):273-9.
55. Jabeen S, Khurshiduzzaman M. Incidence of post obturation pain following single and multi visit root canal treatment in a teaching hospital of Bangladesh. *Mymensingh medical journal : MMJ* 2014;**23**(2):254-60.
56. Liu S, Leng Z-y. Short-term clinical effect of single-visit root-canal therapy in infected root canals. *Journal of Shanghai Jisobong University* 2013;**33**(7):1014-17.

57. Molander A, Warfvinge J, Reit C, Kvist T. Clinical and radiographic evaluation of one- and two-visit endodontic treatment of asymptomatic necrotic teeth with apical periodontitis: a randomized clinical trial. *Journal of endodontics* 2007;**33**(10):1145-8.
58. Mulhern JM, Patterson SS, Newton CW, Ringel AM. Incidence of postoperative pain after one-appointment endodontic treatment of asymptomatic pulpal necross in single-rooted teeth. *Journal of endodontics* 1982;**8**(8):370-5.
59. Oginni A, Udoeye CI. Endodontic flare-ups: comparison of incidence between single and multiple visits procedures in patients attending a Nigerian teaching hospital. *Odonto-stomatologie tropicale = Tropical dental journal* 2004;**27**(108):23-7.
60. Paredes-Vieyra J, Enriquez FJ. Success rate of single- versus two-visit root canal treatment of teeth with apical periodontitis: a randomized controlled trial. *Journal of endodontics* 2012;**38**(9):1164-9.
61. Pekruhn RB. Single-visit endodontic therapy: a preliminary clinical study. *Journal of the American Dental Association (1939)* 1981;**103**(6):875-7.
62. Penesis VA, Fitzgerald PI, Fayad MI, Wenckus CS, BeGole EA, Johnson BR. Outcome of one-visit and two-visit endodontic treatment of necrotic teeth with apical periodontitis: a randomized controlled trial with one-year evaluation. *Journal of endodontics* 2008;**34**(3):251-7.
63. Peters LB, Wesselink PR. Periapical healing of endodontically treated teeth in one and two visits obturated in the presence or absence of detectable microorganisms. *International endodontic journal* 2002;**35**(8):660-7.
64. Prashanth MB, Tavane PN, Abraham S, Chacko L. Comparative evaluation of pain, tenderness and swelling followed by radiographic evaluation of periapical changes at various intervals of time following single and multiple visit endodontic therapy: an in vivo study. *The journal of contemporary dental practice* 2011;**12**(3):187-91.
65. Rao KN, Kandaswamy R, Umashetty G, Rathore VP, Hotkar C, Patil BS. Post-Obturation pain following one-visit and two-visit root canal treatment in necrotic anterior teeth. *Journal of international oral health : JIOH* 2014;**6**(2):28-32.
66. Risso PA, Cunha AJ, Araujo MC, Luiz RR. Postobturation pain and associated factors in adolescent patients undergoing one- and two-visit root canal treatment. *Journal of dentistry* 2008;**36**(11):928-34.
67. Singh S, Garg A. Incidence of post-operative pain after single visit and multiple visit root canal treatment: A randomized controlled trial. *Journal of conservative dentistry : JCD* 2012;**15**(4):323-7.
68. Trope M, Delano EO, Orstavik D. Endodontic treatment of teeth with apical periodontitis: single vs. multivisit treatment. *Journal of endodontics* 1999;**25**(5):345-50.
69. Waltimo T, Trope M, Haapasalo M, Orstavik D. Clinical efficacy of treatment procedures in endodontic infection control and one year follow-up of periapical healing. *Journal of endodontics* 2005;**31**(12):863-6.
70. Wang C, Xu P, Ren L, Dong G, Ye L. Comparison of post-obturation pain experience following one-visit and two-visit root canal treatment on teeth with vital pulps: a randomized controlled trial. *International endodontic journal* 2010;**43**(8):692-7.
71. Weiger R, Rosendahl R, Lost C. Influence of calcium hydroxide intracanal dressings on the prognosis of teeth with endodontically induced periapical lesions. *International endodontic journal* 2000;**33**(3):219-26.
72. Wong AW, Zhang S, Li SK, Zhu X, Zhang C, Chu CH. Incidence of post-obturation pain after single-visit versus multiple-visit non-surgical endodontic treatments. *BMC oral health* 2015;**15**:96.

73. Wong AW, Tsang CS, Zhang S, Li KY, Zhang C, Chu CH. Treatment outcomes of single-visit versus multiple-visit non-surgical endodontic therapy: a randomised clinical trial. *BMC oral health* 2015;**15**(1):162.
74. Yoldas O, Topuz A, Isci AS, Oztunc H. Postoperative pain after endodontic retreatment: single- versus two-visit treatment. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics* 2004;**98**(4):483-7.